arXiv:1201.1212v1 [quant-ph] 5 Jan 2012

Measuring quantumness via anticommutators

Rosario Fazio,[1, 2] Kavan Modi,[3, 2] Saverio Pascazio,[4, 5] Vlatko Vedral,[3, 2, 6] and Kazuya Yuasa[7]
[1] *NEST, Scuola Normale Superiore and Istituto Nanoscienze – CNR, 56126 Pisa, Italy*
[2] *Center for Quantum Technology, National University of Singapore, Singapore*
[3] *Department of Physics, University of Oxford, Clarendon Laboratory, Oxford, OX1 3PU, UK*
[4] *Dipartimento di Fisica and MECENAS, Università di Bari, I-70126 Bari, Italy*
[5] *INFN, Sezione di Bari, I-70126 Bari, Italy*
[6] *Department of Physics, National University of Singapore, Singapore*
[7] *Waseda Institute for Advanced Study, Waseda University, Tokyo 169-8050, Japan*

(Dated: January 6, 2012)

We introduce a method to witness the quantumness of a system. The method relies on the fact that the anticommutator of two classical states is always positive. We show that there is always a nonpositive anticommutator due to any two quantum states. We notice that interference depends on the trace of the anticommutator of two states and it is therefore more natural to detect quantumness by looking at anticommutators of states rather than their commutators.

Introduction. It is usually stated that the main difference between quantum and classical physics is that quantum observables do not commute while all classical properties do. In other words, some properties of quantum systems cannot be specified simultaneously. Well known examples are the position and the momentum of any quantum particle, or the x and z spin components of a spin half particle. However, quantum measurements are only half of the picture. There are also quantum states, which are positive semi-definite, Hermitian operators (density matrices). Classical states, on the other hand, are simply-speaking probability distributions. In other words, all classical states commute with one another, while quantum states in general do not (quantum states, being Hermitian operators, are after all also observables). Can quantumness, therefore, be thought of as the degree of non-commutation of quantum states?

In this Letter we would like to argue that how quantum a system is depends on the non-commutativity of its accessible states. However, it turns out that operationally it is more natural to look at anti-commutators of states rather than their commutators. The reason is that the most basic and natural quantum experiment is to interfere two different states ρ_1 and ρ_2. It is well known [1–3] that the interference fringes depend on the quantity $\operatorname{tr}(\rho_1\rho_2)$, which is the same as half of the trace of the anti-commutator of the two states. The main result of our paper will be that even if two states do not initially anti-commute, it is always possible by repeated measurements of anti-commutators (and only anti-commutators) to reveal any underlying quantumness. Before we show this, we first set the scene by introducing a formal definition of quantumness.

Quantumness. We shall focus for simplicity on finite-dimensional systems and their algebra of observables $\mathcal{A}$ [4, 5]. The following two statements are equivalent [6, 7]: given any pair $X, Y \in \mathcal{A}$,

$$\mathcal{A} \text{ is commutative: } [X,Y] = XY - YX = 0; \qquad (1)$$

$$X \geq 0, Y \geq 0 \longrightarrow \{X,Y\} = XY + YX \geq 0. \qquad (2)$$

Therefore, if the symmetrized product (2) of two positive observables can take negative values, the algebra of observables is non-Abelian and the system is quantum. This motivated the notion of "quantumness witness" [6, 7], that was experimentally tested in [8, 9].

Our objective is to extend the consequences of the characterization (1)-(2) to *states* as well. Let us start by observing that states are non-negative Hermitian operators. Consider a protocol by which an observer $\mathcal{O}$ is given two states ρ_1 and ρ_2, without knowing whether they commute. Assume that $\mathcal{O}$ can only perform interference experiments and therefore can only extract information about *anti*commutators. Can $\mathcal{O}$ understand that the two states do not commute (and therefore that the system is quantum)? Notice that the equivalence between (1) and (2) pertains to the *whole* algebra and not to any given couple of operators. In other words, $[\rho_1, \rho_2] \neq 0$ does not imply in the least that $\{\rho_1, \rho_2\}$ can take negative value. Consider for example two qubits in states $\rho = \frac{1}{2}(\mathbb{1} + \boldsymbol{x}\cdot\boldsymbol{\sigma})$ and $\rho' = \frac{1}{2}(\mathbb{1} + \boldsymbol{x}'\cdot\boldsymbol{\sigma})$. Then $\{\rho, \rho'\}$ is positive definite if $\boldsymbol{x}^2 + \boldsymbol{x}'^2 + \boldsymbol{x}\cdot\boldsymbol{x}' < 1$. For two qubit-states satisfying this condition, the anticommutator (first-order interference experiment) does not bring quantumness to light.

However, $\mathcal{O}$ can perform interference experiments of *any* order, so that $\mathcal{O}$ can in principle obtain information about repeated measurements of any anti-commutators, such as $\{\rho_1, \rho_1\}, \{\rho_1, \rho_2\}, \{\{\rho_1, \rho_1\}, \rho_2\}$ and any (arbitrary) order of nested anticommutators (see Appendix for a general scheme). The main result of this article is a theorem: we shall prove that given any two states of a quantum system, it is possible to bring to light quantumness by *only* looking at the available anticommutators.

Our strategy will be the following. We shall first observe (Theorem 1) that if one of the two states is pure, one anticommutator suffices to bring quantumness to light. Therefore pure states are privileged and presumably quantumness will be easier to detect for states that are "close" to pure states. We shall then observe (Theorem 2) that any mixed state can be made arbitrarily close to a pure state by iterating anticommutators for a finite number of times. The proof is a familiar application of statistical mechanical concepts. Finally, Theorem 3 will show that the anticommutator of any two states that are sufficiently close to pure states is not positive definite. Our result will be valid for qudits of any dimension.

Theorem 1. If a state is pure, $\rho_1 = \rho_1^2 = |\psi\rangle\langle\psi|$, then its anticommutator with any other state ρ_2 is not a non-negative definite operator, provided $[\rho_1, \rho_2] \neq 0$.

Proof. Let $\rho_2 = \sum_i \lambda_i |\phi_i\rangle\langle\phi_i|$, where $\{|\phi_i\rangle\}$ are the eigenvectors of ρ_2 and form a complete orthonormal set of basis vectors, $\sum_i |\phi_i\rangle\langle\phi_i| = \mathbb{1}$, and $\{\lambda_i\}$ are the eigenvalues of ρ_2, satisfying $0 \leq \lambda_i \leq 1$ and $\sum_i \lambda_i = 1$. Its anticommutator with the pure state ρ_1 yields

$$\{\rho_1, \rho_2\} = \sum_i \lambda_i (f_i |\psi\rangle\langle\phi_i| + f_i^* |\phi_i\rangle\langle\psi|), \tag{3}$$

where $f_i = \langle\psi|\phi_i\rangle$ and $\sum_i |f_i|^2 = \sum_i \langle\psi|\phi_i\rangle\langle\phi_i|\psi\rangle = 1$. We normalize this anticommutator as

$$\rho_{12} = \frac{\{\rho_1, \rho_2\}}{\mathrm{tr}\{\rho_1, \rho_2\}} = \frac{\sum_i \lambda_i (f_i |\psi\rangle\langle\phi_i| + f_i^* |\phi_i\rangle\langle\psi|)}{2\sum_i \lambda_i |f_i|^2}. \tag{4}$$

Since ρ_{12} is a Hermitian and unit-trace operator, its purity $\mathrm{tr}\rho_{12}^2$ must be less than 1 to satisfy positivity condition. If on the other hand purity exceeds unity, ρ_{12} is proved not to be non-negative definite. We get

$$\mathrm{tr}\rho_{12}^2 = \frac{(\sum_i \lambda_i |f_i|^2)^2 + \sum_i \lambda_i^2 |f_i|^2}{2(\sum_i \lambda_i |f_i|^2)^2}, \tag{5}$$

and hence, positivity of $\{\rho_1, \rho_2\}$ is violated if $\mathrm{tr}\rho_{12}^2 > 1$, namely, if

$$\sum_i \lambda_i^2 |f_i|^2 > \left(\sum_i \lambda_i |f_i|^2\right)^2. \tag{6}$$

This is always true by virtue of the Cauchy-Schwarz inequality (recall that $\sum_i |f_i|^2 = 1$), except when $\lambda_i f_i \propto f_i$ for all i, i.e., except when $\mathrm{tr}\rho_{12}^2 = 1$. On the other hand, $\mathrm{tr}\rho_{12}^2 = 1$ implies

$$\begin{aligned}[\rho_1, \rho_2] &= \sum_i \lambda_i (f_i |\psi\rangle\langle\phi_i| - f_i^* |\phi_i\rangle\langle\psi|) \\ &\propto \sum_i (f_i |\psi\rangle\langle\phi_i| - f_i^* |\phi_i\rangle\langle\psi|) = 0.\end{aligned} \tag{7}$$

This proves the theorem. □

Theorem 2. If a density operator ρ does not have a degeneracy in its maximum eigenvalue, then the normalized operator corresponding to ρ^n approaches, as $n \to \infty$, the pure state of the eigenvector corresponding to the largest eigenvalue.

Proof. Let us write state ρ in its eigenbasis,

$$\rho = \sum_i \lambda_i |i\rangle\langle i|, \tag{8}$$

λ_0 being the largest eigenvalue, with no loss of generality. Since $\lambda_i \leq 1$ in general, λ_i^n decays as $n \to \infty$, but λ_0^n decays most slowly, and we end up with

$$\lim_{n\to\infty} \frac{\rho^n}{\mathrm{tr}\rho^n} = \lim_{n\to\infty} \sum_i \frac{\lambda_i^n}{\sum_j \lambda_j^n} |i\rangle\langle i| = |0\rangle\langle 0|, \tag{9}$$

where $|0\rangle$ is the eigenstate belonging to the (non-degenerate) largest eigenvalue. □

Instead of taking n to infinity and bringing a given mixed state to a pure state, like in Theorem 2, we may just take a finite number of iterations and bring the state close to the pure state. Given any two mixed states, we can take them both ϵ-close to pure states, respectively. The anticommutator of these two states will admit (at least) a negative eigenvalue. Below we bound ϵ based on how close these two states are to each other.

Observe first that we can define the closest pure state $|\psi\rangle$ to an arbitrary state ρ, in the sense that $\langle\psi|\rho|\psi\rangle$ is maximum among all pure states, and ρ can be expressed as a convex sum

$$\rho = \lambda |\psi\rangle\langle\psi| + (1-\lambda)\sigma, \tag{10}$$

with a density operator σ ($\sigma \geq 0$, $\mathrm{tr}\sigma = 1$) that is orthogonal to $|\psi\rangle$, i.e., $\sigma|\psi\rangle = 0$ (λ being the largest eigenvalue of ρ). We are ready to prove our central result.

Theorem 3. Given two mixed states ρ_1 and ρ_2 close to pure states $|\psi_1\rangle$ and $|\psi_2\rangle$, respectively, i.e.,

$$\rho_i = (1-\epsilon_i)|\psi_i\rangle\langle\psi_i| + \epsilon_i \sigma_i \quad (i = 1, 2) \tag{11}$$

with $\sigma_1|\psi_1\rangle = \sigma_2|\psi_2\rangle = 0$, their anticommutator $\{\rho_1, \rho_2\} \ngeq 0$, provided ϵ_1 and ϵ_2 are small enough to satisfy

$$\epsilon_1 g_1 + \epsilon_2 g_2 < (1 - |f|^2)/2, \tag{12}$$

where $g_1 = \langle\psi_2|\sigma_1|\psi_2\rangle$, $g_2 = \langle\psi_1|\sigma_2|\psi_1\rangle$ and $f = \langle\psi_1|\psi_2\rangle$.

Proof. The anticommutator of ρ_1 and ρ_2 reads

$$\begin{aligned}\{\rho_1, \rho_2\} =& (1-\epsilon_1)(1-\epsilon_2)(f|\psi_1\rangle\langle\psi_2| + f^*|\psi_2\rangle\langle\psi_1|) \\ &+ \epsilon_1(1-\epsilon_2)(\sigma_1|\psi_2\rangle\langle\psi_2| + |\psi_2\rangle\langle\psi_2|\sigma_1) \\ &+ \epsilon_2(1-\epsilon_1)(\sigma_2|\psi_1\rangle\langle\psi_1| + |\psi_1\rangle\langle\psi_1|\sigma_2) \\ &+ \epsilon_1\epsilon_2\{\sigma_1, \sigma_2\},\end{aligned} \tag{13}$$

with its trace given by

$$\begin{aligned}\mathrm{tr}\{\rho_1, \rho_2\} =& 2(1-\epsilon_1)(1-\epsilon_2)|f|^2 \\ &+ 2\epsilon_1(1-\epsilon_2)g_1 + 2\epsilon_2(1-\epsilon_1)g_2.\end{aligned} \tag{14}$$

Like in Theorem 1, the anticommutator is not positive-definite, $\{\rho_1, \rho_2\} \ngeq 0$, if the purity of $\rho_{12} = \{\rho_1, \rho_2\}/\mathrm{tr}\{\rho_1, \rho_2\}$ exceeds unity. This is readily calculated, at first order in ϵ_1 and ϵ_2:

$$\mathrm{tr}\rho_{12}^2 = \frac{(1 - 2\epsilon_1 - 2\epsilon_2)(1 + |f|^2) + 2\epsilon_1 g_1 + 2\epsilon_2 g_2}{2[(1 - 2\epsilon_1 - 2\epsilon_2)|f|^2 + 2\epsilon_1 g_1 + 2\epsilon_2 g_2]}, \quad (15)$$

and we get $\mathrm{tr}\rho_{12}^2 > 1$ under condition (12). □

Notice that condition (12) becomes more and more stringent when $|f|^2$ is closer to unity, namely, when the two states $|\psi_1\rangle$ and $|\psi_2\rangle$ are very close to each other. Let us also observe that if the maximum eigenvalue is degenerate, the procedure outlined here (that hinges upon nested anticommutators) does not work, simply because it is unable to unearth the negativity of the "final" anticommutator. This is easily understood by looking at Eqs. (10)-(11): the limiting state of Theorem 2 would not necessarily be close to the boundary.

We are now in position to put the results above in perspective. We have given a method to test for quantumness of a system that is preparable in two noncommuting states. However, depending on the purity of these states, we may require many copies to reveal the quantum feature, namely the nonpositivity of the anticommutator. Operationally, given states ρ_1 and ρ_2 we may have to purify them in some finite rounds. To do so we consider the anticommutator of each state with itself, which will purify the state due to Theorem 2. Once we attain desired purification, we can take the anticommutator of the two purified states. Mathematically we can say that given two noncommuting states ρ_1 and ρ_2 we have an nonpositive operator corresponding to a nested anticommutator:

$$\Big\{ \{\rho_1, \dots \{\rho_1, \rho_1\}\}, \{\rho_2, \dots \{\rho_2, \rho_2\}\} \Big\} \propto \{\rho_1^m, \rho_2^n\}. \quad (16)$$

The positivity of the operator on the left and on the right are the same. The equality is lacking only due to different normalization on the two sides. The values of m and n can be interpreted as the number of copies of the states ρ_1 and ρ_2 respectively, that are needed to witness the quantum feature of the system. Note that $m = n = 1$ when one of the states is pure due to Theorem 1. Conversely, note that for highly mixed states m and n take large values to witness quantumness, which is contrary to the Bohr correspondence principal, which asserts that in the large number limit quantum goes to classical. For a related study with on this topic see [13, 14].

Quantum discord. Now we show that the notion of quantumness (of a single system) presented in this Letter is related to quantum discord. Quantum discord and related measures [10] attempt to quantify the quantum correlations in multipartite quantum states. However, for simplicity we only work with bipartite states ρ_{AB} here. Let us denote quantum discord in ρ_{AB} as $D(A|B)$. The notation implies that discord is measured by B, while $D(B|A)$ would imply discord as measured by A. Note that $D(A|B) \neq D(B|A)$.

Suppose Alice and Bob share a state and Alice has to convince Bob that he is quantum correlated to her. She can do this by making two measurement of her system such that the corresponding conditional states of B do not commute. Then she simply has to communicate the outcomes of her measurements and Bob can carry out the interference experiment, i.e. measure the nonpositivity of the anticommutator of the two states prepared by Alice. However, Alice can only prepare conditional states of Bob that do not commute if he is quantum correlated to her. This comes from the following theorem.

Theorem 4. [Chen et al. [11]] A bipartite state is quantum correlated for B if and only if for any set of location operation by A, the conditional state of B all commute:

$$D(A|B) = 0 \quad \Longleftrightarrow \quad \left[\rho_{B|i}\ ,\ \rho_{B|j}\right] = 0 \quad \forall\ i, j, \quad (17)$$

where $\rho_{B|i} = \mathrm{tr}_A[\Lambda_A^i \otimes \mathcal{I}_B(\rho_{AB})]$ are the conditional states of B for a (generalized) quantum operations Λ_A^i that is made on A. Conversely, if $D(A|B) > 0$, then A can make two local operations yielding two conditional states for B that do not commute.

Once Alice remotely prepares two states for Bob, he can measure the anticommutator of these states. In other words, carry out the procedure of Theorems 1-3 above. The anticommutator will be nonpositive if and only if Bob was quantum correlated to Alice. However the number of states that Alice has to produce for Bob depends on the mixedness of the conditional states and how noncommuting they are.

Conclusions. Separating the quantum world from the classical world is an old problem. Normally, one takes a classical limit, i.e. $\hbar \to 0$ or $n \to \infty$ to attain classical mechanics. In this Letter we have introduced a method to witness the quantumness of a system that is preparable in noncommuting states. The method relies on the fact that the anticommutator of two classical states is always positive. We show that there is always a nonpositive anticommutator due to any two noncommuting states. However, the positivity of the anticommutator is dependent on purity of the states. For highly mixed states we require many copies of the two states (or alternatively high-order interference) in order to witness quantumness. In other words quantumness emerges in the large number limit. On the other hand even though quantumness grows with the number of states, detecting the witness remains difficult. This is because when working with n states, we need n sets of n states to witness quantumness (see Appendix). Having n copies of n states may be even reasonable, however, (coherently) interacting all of these states may not be (the coherent interaction is the controlled-SHIFT operator in Fig. 1). Therefore, in the end, the scheme presented here is in agreement with our observation of the world. It says that a macroscopic



Figure 1. H is the Hadamard gate, S is the SHIFT gate. The expectation value of the control qubit (top qubit) in z–basis is the trace of the product $\mathrm{tr}[\rho_1\rho_2\rho_3\rho_4]$.

object is quantum and one can even witness this quantumness, provided enough copies and apparatus that can (coherently) interact these objects. According to this scheme the lack of quantumness in the macroscopic world is due to the limitations on coherent interactions of large number of macroscopic systems. Lastly, we have linked this notion of quantumness to quantum correlation as quantified by quantum discord.

KM and VV acknowledge the support of the John Templeton Foundation. KM, RF, and VV acknowledge the financial support by the National Research Foundation and the Ministry of Education of Singapore. SP and KY would like to thank the Centre for Quantum Technologies of National University of Singapore for the hospitality. SP and KY are partially supported by the Joint Italian-Japanese Laboratory on "Quantum Technologies: Information, Communication and Computation" of the Italian Ministry for Foreign Affairs. KY is supported by the Ministry of Education, Culture, Sports, Science and Technology, Japan.

APPENDIX. *Measuring anticommutator.* Experimentally measuring the positivity of the anticommutator if fully doable. Let us begin with the trace of the anticommutator: $\mathrm{tr}[\mathcal{O}] = \mathrm{tr}[\rho_1\rho_2 + \rho_2\rho_1] = 2\mathrm{tr}[\rho_1\rho_2] = \sum_i \lambda_i$, where λ_i are the eigenvalues of $\mathcal{O}$. If the quantum system is of d-dimensions then there are d eigenvalues and if we know the values of $\mathrm{tr}[\mathcal{O}^l] = \sum_i \lambda_i^l$ for all values of $1 \leq l \leq d$, then we can algebraically solve for $\{\lambda_i\}$. To find the eigenvalues of the $\mathcal{O}$ we have to measure $j_l = \mathrm{tr}[(\rho_1\rho_2)^l]$.

To measure the product of two operators in an experimental setting we make use of the controlled-SHIFT operator [1, 15, 16]. The SHIFT operator's action is defined as: $S\,|\psi_1, \psi_2, \ldots, \psi_{l-1}, \psi_l\rangle = |\psi_l, \psi_1, \psi_2, \ldots, \psi_{l-1}\rangle$. The trace of the SHIFT operator's action from one side only yields

$$\mathrm{tr}[S(\rho_1 \otimes \rho_2 \otimes \cdots \otimes \rho_l)] = \mathrm{tr}[\rho_1\rho_2 \cdots \rho_l]. \tag{18}$$

Notice the tensor product on the l.h.s. and the ordinary matrix product on the r.h.s.

Using a control qubit and implementing the controlled-SHIFT operator, $\mathcal{C}_S = |0\rangle\langle 0| \otimes \mathbb{I} + |1\rangle\langle 1| \otimes S$, we can measure the trace of the product of any number of operators, see Fig. 1 for a circuit representation. Measuring the control qubit in z–basis will give the product of the density operators. However to measure the lth power we will need l copies of the density operator and a suitable SHIFT operation. These resources may not be trivially available and therefore witnessing the quantumness may prove to be difficult.

[1] A. K. Ekert, C. M. Alves, D. K. L. Oi, M. Horodecki, P. Horodecki and L. C. Kwek, Phys. Rev. Lett. **88**, 217901 (2002).

[2] H. A. Carteret, Phys. Rev. Lett. **94**, 040502 (2005).

[3] E. Sjoqvist, A. K. Pati, A. Ekert, J. S. Anandan, M. Ericsson, D. K. L. Oi, and V. Vedral, Phys. Rev. Lett. **85**, 2845 (2000).

[4] H. Araki, *Mathematical Theory of Quantum Fields* (Oxford, Oxford University Press, 2000).

[5] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics* vols 1-2 2nd edn (Berlin, Springer, 2002).

[6] R. Alicki and N. Van Ryn, J. Phys. A: Math. Theor. **41** 062001 (2008).

[7] R. Alicki, M. Piani, and N. Van Ryn, J. Phys. A: Math. Theor. **41**, 495303 (2008).

[8] G. Brida, I. P. Degiovanni, M. Genovese, V. Schettini, S. V. Polyakov, and A. Migdall, Optics Express **16**, 11750 (2008).

[9] G. Brida, I. P. Degiovanni, M. Genovese, F. Piacentini, V. Schettini, N. Gisin, S. V. Polyakov, and A. Migdall, Phys. Rev. A **79**, 044102 (2009).

[10] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, arXiv:1112.6238 [quant-ph].

[11] L. Chen, E. Chitambar, K. Modi, and G. Vacanti, Phys. Rev. A **83**, 020101 (2011).

[12] A. K. Ekert, C. M. Alves, D. K. L. Oi, M. Horodecki, P. Horodecki, and L. C. Kwek, Phys. Rev. Lett., **88**, 217901 (2002).

[13] K. Modi, R. Fazio, S. Pascazio, V. Vedral and K. Yuasa, "Classical to quantum in large number limit" arXiv:1112.5830 [quant-ph].

[14] P. Facchi, S. Pascazio, V. Vedral and K. Yuasa "Quantumness and Entanglement Witnesses" arXiv:1111.5257 [math-ph].

[15] M. Keyl and R. F. Werner, Phys. Rev. A **64**, 052311 (2001).

[16] V. Vedral, Introduction to quantum information science. Oxford: Oxford University Press (2007).